

Comparative Fingerprint and Extraction Yield of Medicinal Herb Phenolics with Hepatoprotective Potential, as Determined by UV-Vis and FT-MIR Spectroscopy

Simona ZAVOI¹, Florinela FETEA¹, Floricuta RANGA¹, Raluca M. POP¹, Anca BACIU², Carmen SOCACIU^{1*}

¹University of Agricultural Sciences and Veterinary Medicine, Faculty of Agriculture, 3-5 Mănăstur Street, Cluj-Napoca, Romania; carmen.socaciu@usamvcluj.ro (*corresponding author)

²Center for Applied Biotechnology CCD-BIODIATEC, Proplanta Cluj-Napoca, Romania

Abstract

The present study was aimed to compare the polyphenolic composition of six medicinal herbs, from wild flora of Romania. The plants investigated, *Cynara scolimus* (artichoke), *Taraxacum officinalis* (dandelion), *Chelidonium majus* (celandine), *Hypericum perforatum* (St. John's wort), *Silybum marianum* (Mary thistle) and *Lycopodium clavatum* (Wolf's claw) are known, to have hepatoprotective action. Using in parallel glycerol-water, ethanol-water and methanol, the solvent-dependence of the extract fingerprint and composition in bioactive molecules was studied by UV-Vis and Infrared (FT-MIR) spectrometry. The extraction yields, calculated as an extraction factor (EF) were superior in acidic methanol comparative to glycerin and ethanol, favorising the increase in phenolic acids against flavonoid derivatives. Based on the differences of polarity between the three solvents used, higher EF values were obtained for dandelion, artichoke, celandine and St. John wort, more rich in phenolic acids than flavonoids. Mary thistle and Wolf's claw had lower concentrations of phenolics, but higher content of lignans and terpenoids. Based on the FT-MIR peaks from 8 regions, for each plant extract, has been determined the fingerprint region between 900 and 1500 cm⁻¹ and identified the specific functional groups. A good, significant correlation was found between the concentration of total phenolics calculated by UV-Vis spectrometry and FTIR methods, after calibration with gallic acid. The value of the MIR signal at 1743 cm⁻¹ may be considered a good indicator of phenolics concentration in such extracts. Combined UV-Vis and FTIR spectroscopy are recommended as rapid and reliable tools to investigate the fingerprint and to predict the composition of medicinal plants or to evaluate the quality and authenticity of different standardized formulas.

Keywords: FT-MIR, hepatoprotection, medicinal herbs, quality and authenticity, UV spectrometry

Introduction

The traditional herb medicines showed, since centuries, beneficial effects on health promotion, out of side effects, as compared with synthetic drugs. It is also known that their composition is dependent on many ontogenic or genotypic factors influenced by their environment, age, time of harvesting, drying and storage, as well the solvent used to obtain extracts. Due to their natural heterogeneity, the quality of herbs from wild environments shows great fluctuations, so their standardization has been extensively promoted during the last years, the following three attributes being verified: authenticity, purity and assay of their action (Hussain *et al.*, 2009; Yadav and Dixit, 2008). The identification of phytochemicals' fingerprint by chromatography and spectroscopy may provide effective information about qualitative and quantitative composition of herbal medicines and their pattern recognition can be achieved by chemometry (Bender, 2005; Maloney, 2004) and used to discriminate among different herbs and extracts, with

noticeable role in the development of standardized formulas, like other pharmaceuticals to make these remedies evidence-based medicines, (Giri *et al.*, 2010; Liang *et al.*, 2004; McGuffin *et al.*, 1997; Yadav and Dixit, 2008), according to WHO requirements (WHO, 2003)

The evaluation of a herbal product by its metabolomic fingerprinting can be accomplished by appropriate methods, including HPLC with UV (DAD), ELSD, MS detection or GC-MS, HPTLC-densitometry, FT-MIR, NIR, NMR or a combination of these techniques (Fan *et al.*, 2006; Giri *et al.*, 2010; Gong *et al.*, 2006; 2009; Hashimoto and Kameoka, 2008; Li *et al.*, 2008; Mattoli *et al.*, 2006).

The UV-Vis spectroscopy offer a simple, cheap and easy-to-use technique to identify and quantify the main phytochemicals, discriminating between the lipophilic and hydrophilic phytochemicals, in relation to the polarity of the extraction solvent.

Fourier transform infrared spectroscopy (FTIR) offers a rapid and non-destructive investigation, easy to

use to fingerprint herbal extracts or powders, is relatively uncommon compared with chromatographic and classical methods (Hussain *et al.*, 2009a; Li *et al.*, 2004b; Liu *et al.*, 2006). The use of attenuated total reflectance (ATR) device evolved rapid FTIR measurements of liquids such as oils and plant extracts, allowing the identification and quantification of valuable plant biomarkers (Schultz and Baranska, 2007).

The herbs, known traditionally, to have hepatoprotective action are generally rich in polyphenols with high antioxidant potential (Mary thistle, artichoke, dandelion, greater celandine, St. John's wort, etc.) since major liver diseases are related to oxidative stress and cellular necrosis (Negi *et al.*, 2008; Utrilla, 1996).

Dandelion (*Taraxacum officinalis*) is an old remedy for glandular unbalances, used in the therapy of liver diseases for the biliary stimulation, normalization of blood circulation, diuresis and toxins release. Its bitter substances (eudesmanolides and germacranolides) named generically "taraxacin", are anti-vomitives and antioxidants (Jeon *et al.*, 2008). Dandelion contains as well flavonoids (7-D glucosides of apigenol and luteoline) and steroids (stigmasterol, sitosterol, ergosterol) which block the *de novo* synthesis of cholesterol and are competitors of cholesterol deposition (Williams *et al.*, 1996). It contains also inulin with protective action against diabetes and liver diseases (Schütz *et al.*, 2006), similarly to artichoke.

Since many years it was reported that artichoke (*Cynara scolimus*) induces regeneration of rat liver (Maros *et al.*, 1966), mainly due to cynarin, a phenolic derivative which stimulates liver cell excretion (Li *et al.*, 2004a; Wang *et al.*, 2003). Luteolin, a flavone which inhibit the *de novo* cholesterol synthesis as well sesquiterpene lactones has coleretic and cholagogue effects (Saenz Rodriguez *et al.*, 2002). The leaf extract of artichoke reduces mild dyspepsia (Marakis *et al.*, 2002) and have antioxidant potential, as demonstrated on leukocytes (Perez-Garcia *et al.*, 2000).

Greater celandine (*Chelidonium majus*) is a medicinal poppy (Bone, 1996) with beneficial antihepatotoxic and meanwhile controversial effects on liver (Duke, 1985; Mitra *et al.*, 1992; 1996). It is rich in alkaloids (Gu *et al.*, 2010), mainly chelidonin, a naphthofenantridin derivative which stimulates the enzymatic production in liver and pancreas, it is colecystokinetic and antispastic (Gilca *et al.*, 2010; Hriscu *et al.*, 1980). It was demonstrated experimentally to be antiinflammator, anticancerigenic și antimicrobial (Vavreckova *et al.*, 1996) being recommended in cyrosis and chronic hepatitis therapy. Using a standardized extract (4 mg chelidonin for 6 weeks) for patients with digestive syndroms, a significant reduction of symptoms was reported (Ritter and Schatton, 1993). Sanguinarine is another alkaloid found in celandine which acts as a colchicine-like cytostatic (Lopus and Panda, 2006; Malikova *et al.*, 2006) with antiviral, antiinflammatory and antibacterial potential (Zdařilová *et al.*, 2006). Meanwhile, overdoses

are recommended to avoid, due to its alkaloid charge and hepatotoxicity (Benninger *et al.*, 1999).

Mary thistle (*Silybum marianum*) is known to have hepatoprotective action and to be a hepatocyte activator (Ferenci *et al.*, 1989; Salmi and Sarna, 1982), being used not only as tea but also in some standardized drugs such as Silimarine (generic name of lignan flavonoids, a mixture of silibine, silibinin, silicristine and silidianine). Silibine is the most active derivative, the extracts being standardized at 70-80% silibine. The mechanisms involved in its hepatoprotective action are diverse and include antioxidant and antiperoxidation effects (Basaga *et al.*, 1997; Bosisio *et al.*, 1992), detoxifying effect (Baer-Dubowska *et al.*, 1998; Miguez *et al.*, 1994), antiviral action (McPartland, 1996) and glutathion protection (Cabrera, 1996). Silimarine protects hepatocytes against toxic effects of acetaminophen, ethanol, carbon tetrachlorine (Bosisio *et al.*, 1992; Favari and Perez-Alvarez, 1997; Muriel *et al.*, 1992), decrease the fibrosis by III-peptide procollagen inhibition (Feher *et al.*, 1989) and inhibit cytochrome P 450 (Baer-Dubowska *et al.*, 1998).

St. John's wort (*Hypericum perforatum*) contains several compounds with hepatoprotective properties, by synergistic action of hypericins and phenolics. Hypericins are reddish-violet condensed anthraquinones found (0,1-0,2%) in leaves and flowers. Due to their UV fluorescence, are photoactivated in sun, being good candidates for tumor photodynamic therapy. It has also anti-depressive and antiviral, has cholagogue and choleretic properties (Ali and Olivo, 2002; Kurth and Spreeman, 1998). Flavonoid glucosides (rutoside) protects the liver against oxidative stress, especially by its action on superoxid dismutase (Istodor, 2003). The condensed tannins and hyperforin, have antibiotic effects, inhibits cytochrom P450 CYP3A4 și CYP2C9 (Barnes *et al.*, 2001; Gitea *et al.*, 2007; Kurth and Spreeman, 1998; Nahrstedt and Butterweck, 1997).

Wolf's claw (*Lycopodium clavatum*) is a less studied herb, which contains more than 35 alkaloids and other biomolecules, whose structure and pharmacologic effects are not yet elucidated. The bioactive molecules are lycopodine, clavatoxin, clavolonin, clavatin, liclanitin, annapodin, izolzopodin, as well nicotine, triterpenoids and sterols, flavonoids, radium (known as an antitumor mineral). Triterpenoids and sterols are stimulators of digestive system, antialergic, prevent the cholesterol and fats deposition, may have good effects on liver tumor inhibition, anti-inflammatory and antimicrobial action (Betancor-Fernandez *et al.*, 2003).

The present study aimed to compare the fingerprint of different extracts of the above-mentioned medicinal herbs, collected from wild flora of Romania. The dependence of the extract composition on the solvent polarity (glycerol-water *vs* ethanol-water *vs* acidic methanol) was studied by UV-Vis spectrometry and Infrared (FT-MIR) fingerprints.

Materials and methods

Medicinal plants and preparation of extracts

Six types of medicinal plants, from wild flora of different areas of Transylvania, Romania were investigated. The plants were numbered as follows: 1-*Cynara scolimus* (artichoke), 2-*Taraxacum officinalis* (dandelion), 3-*Chelidonium majus* (celandine), 4-*Hypericum perforatum* (St. John's wort), 5-*Silybum marianum* (Mary thistle), 6-*Lycopodium clavatum* (Wolf's claw). Aliquots of 15 g from each dried and grounded plant (selected from 100 g mix of leaves, stems and flowers) were extracted in 85 ml solvent: methanol 90% in water, acidulated with 1% hydrochloric acid (M) or ethanol 93% in water (E) or 30% glycerin in water (G). After sonication 30 min, centrifugation and filtration, the clear extracts were kept in the deep freezer until analysis.

UV-Vis spectra and calculation of extraction factors

The UV-Vis spectra were recorded (700-200 nm) for each extract (M, E or G) using a Jasco V 530 Spectrophotometer. There were identified the maxima wavelengths specific for phenolics (280 and 330 nm), carotenoids (420-470 nm) and/or chlorophylls (663 nm).

To compare the yields of extraction in different solvents it has been calculated the Extraction Factors (EF) of bioactive molecules from each extract, considering the ab-

sorption values ($A_{\lambda_{\max}}$) recorded for each $\lambda_{\max}$, multiplied with the dilution factor (d). The formula applied was:

$$EF = A(\lambda_{\max}) \times d$$

The results were expressed as mean values of four samples per plant and in duplicated extracts from each plant. The content of total phenolics was determined by Vis spectrometry, using the standard method, Folin Ciocalteu (Singleton, 1999)

FT-MIR measurements

The Fourier Transform Infrared spectrum (FTIR) of each extract was recorded in the MIR region, from 4000 to 900 cm^{-1} , and 64 scans were accumulated for each spectrum using the Horizontal Attenuated Total Reflection (HATR) device, using a Shimadzu Prestige 2 FTIR spectrometer (with apodization Happ-Genzel). The spectral data were processed using the IR solution Software Overview (Shimadzu) and OriginR 7SR1 Software (Origin-Lab Corporation, Northampton, USA). The spectra were registered as fluid (M, E, G) but also evaporated extracts (these data are not shown). Total phenolics were determined also by FTIR method, either using the intensity of the peak at 1742 cm^{-1} or from the area of the region 950-1900 cm^{-1} , considering the calibration curve with pure gallic acid (range of concentrations 5 to 30 mg/ml methanol).

Results and discussions

Extraction factors of bioactive molecules, based on UV-Vis spectra

The comparative UV-Vis spectra of the ethanol (E) and glycerin (G) extracts of the six medicinal plants were recorded (data not shown), as well in methanolic extracts (M) (Fig. 1), methanol being considered a "reference" solvent known to extract phenolics and terpenoids from these plants.

Based on their specific spectra, the absorption maxima ($\lambda_{\max}$) of each plant extract and the mean values of extraction factors (EF) were calculated for each solvent (E, G and M) (Tab. 1).

To have an integrated image of the differences between plants, solvent type and concentrations of bioactive molecules extracted, the EF mean values at 270-290 nm (for phenolic acid derivatives extracted in E, G, M) (EFE1, EFG1, EFM1) and at 317-340 nm (for flavonoid derivatives) (EFE2, EFG2, EFM2) for each of the 6 plants were represented (Fig. 2).

According to Tab. 1 and Fig. 2 data, it has been noticed that extraction factors in acidic methanol were superior to glycerin and ethanol, especially for phenolic acids (EFE1, EFG1, EFM1) comparing to flavonoid derivatives (EFE2, EFG2, EFM2). Based on the differences of polarity between the three solvents used (glycerin the most polar, followed by methanol and ethanol) it has been noticed high-

Tab. 1. The absorption maxima $\lambda_{\max}$ (nm) of each plant extract from UV-Vis spectra and the mean values calculated ($X \pm SD$) for extraction factors (EF)

Plant (nr.)	$\lambda_{\max}$ (nm)	EFE Ethanol	EFG Glycerin	EFM Methanol
Dandelion (1)	279	7.69 ± 0.55	231.64 ± 10.50	240 ± 0.55
	320	7.96 ± 0.58	201.24 ± 9.20	280 ± 10.50
	396	2.07 ± 0.12	106.09 ± 7.50	0
	247	83.08 ± 3.08	43.85 ± 1.50	0
Artichoke (2)	295	82.94 ± 3.88	41.19 ± 1.35	91.71 ± 6.50
	327	97.05 ± 5.02	42.86 ± 1.65	73.31 ± 2.80
	663	0	0	6.03
	212	0	255.36 ± 11.34	0
celandine (3)	275	67.79 ± 2.05	103.96 ± 6.90	288 ± 12.10
	330	48.94 ± 1.86	72.96 ± 2.50	219 ± 10.13
	396	13.14 ± 0.90	42.86 ± 1.98	0
	663	0	0	9.81 ± 0.40
St. John's wort (4)	270	33.89 ± 1.80	137.71 ± 8.98	422 ± 21.05
	330	19.03 ± 1.10	82.68 ± 3.98	238 ± 11.08
Mary thistle (5)	280	19.76 ± 1.06	34.88 ± 1.98	142 ± 9.58
	325	14.63 ± 1.01	29.26 ± 1.27	77 ± 3.10
	395	19.76 ± 1.12	18.01 ± 1.18	0
Wolf's claw (6)	233	57.91 ± 2.90	0	108.31 ± 9.80
	284	31.92 ± 1.75	25.81 ± 1.39	0
	324	29.94 ± 1.50	24.96 ± 1.35	59.4 ± 2.80
	397	12.46 ± 1.80	3.99 ± 0.84	0
	652	0	0	2.69 ± 0.54




Fig. 1. Comparative UV-Vis spectra of methanolic (M) extracts of *Taraxacum officinalis* (1); *Cynara scolimus* (2); *Chelidonium majus* (3); *Hypericum perforatum* (4); *Silybum marianum* (5); *Lycopodium clavatum* (6)




Fig. 2. Comparative mean values of extraction factors (EF) in different solvents (ethanol-E, glycerin-G, methanol-M) at 270-290 nm (EFE1, EFG1, EFM1) and 317-340 nm (EFE2, EFG2, EFM2) for each of the studied plants (series 1-6)

er EF values for dandelion (1), celandine (3) and St. John wort (4), more rich in polar molecules, such as phenolic acids and flavonoids. Plants 1 and 3 had similar EF in M and G, plant 2 (artichoke) was better extracted in ethanol and was more rich in phenolic acid derivatives. St. John's wort (4) components were two times better extracted in methanol than glycerin, and low EF values in ethanol, an indication of polar active molecules. Mary thistle (5) and Wolf's claw (6) contained reduced concentrations of phenolics, but high absorptions in methanol at 280 and 233 nm, respectively, which might be attributed to higher concentrations of lignans and terpenoids.

For therapeutic reasons it has been considered that ethanol extracts or evaporated methanolic extracts can provide higher concentrations of bioactive molecules from these plants. Anyway, as it was reported recently, methanolic extracts showed hepatoprotective activity against Carbon Tetrachloride-Induced Hepatotoxicity in Rats (Ahsan *et al.*, 2009). Of course for humans the total elimination of methanol (under vacuum) is a condition to have a safe standardized extract.

Tab. 2. Absorption peak areas of different regions (1-8) of FT-MIR spectra recorded for methanol extracts (M) of the studied medicinal plants

Wavenumber (1/cm)	1	2	3	4	5	6
3300-3350 (8)	0.70	0.92	0.63	0.82	0.93	0.80
2800-3000 (7)	0.89	0.81	0.98	0.90	0.82	0.75
1500-1520 (5)	0.35	0.22	0.30	0.52	0.32	0.30
1300-1440 (4)	0.72	0.80	0.80	0.95	0.43	0.63
1170-1230 (3)	0.90	0.90	0.92	1.15	0.78	0.80
1000-1100 (2)	1.35	1.45	1.32	1.50	1.05	1.28
<1000 (1)	0.50	0.78	0.60	0.85	0.60	0.52
Total phenolics (by FTIR)	11.33	11.03	11.16	10.18	8.05	9.26
mg GA eq./ml M						
Total phenolics (by Vis spectrometry)	9.307	9.066	9.171	8.386	6.669	7.642
mg/ GA eq./ml M						


FT-MIR fingerprint

The FT-MIR spectra (4000-900 cm^{-1}) of E and G extracts of each plant were registered and the specific wavenumbers and intensities were considered (data not shown). Fig. 3 presents the FT-MIR spectra of M extracts and Tab. 2 includes the corresponding absorption peak areas for specific regions (1-8). In Tab. 2, is included the total phenolics concentration in methanol (M) extracts determined by FTIR and by Vis spectrometry.


The functional groups identification was based on the FTIR peaks attributed to stretching and bending vibrations. Eight areas (marked from 1 to 8) (Fig. 3) were identified in the MIR domain and the fingerprint region was localized between 900 and 1500 cm^{-1} (areas 1-4).

Area 1 (< 1000 cm^{-1}) corresponds to C-H bending vibrations from isoprenoids, area 2 (997-1130 cm^{-1}) to stretching vibrations C-O of mono-, oligo- and carbohydrates, with signals at 1030, 1054, 1104, and 1130 cm^{-1} , while area 3 (1150-1270 cm^{-1}) corresponds to stretch-


1




4




2



5



3



6




Fig. 3. The FTIR fingerprint of the metanolic (M) extracts of the studied plants: *Cynara scolimus* (1); *Taraxacum officinalis* (2); *Chelidonium majus* (3); *Silybum marianum* (4); *Hypericum perforatum* (5); *Lycodium clavatum* (6). The specific regions are numbered 1 to 8




Fig. 4. Comparative representation of the differences recorded between the areas of regions 1, 2, 4, 6, 7 corresponding to each plant extract (ethanol-E, glycerin-G or methanol-M)

ing vibrations of carbonyl C-O or O-H bendings. Area 4 ($1300-1450\text{ cm}^{-1}$) correspond to stretching vibrations C-O (amide) and C-C stretchings from phenyl groups, while area 5 ($1500-1600\text{ cm}^{-1}$) to aromatic domain and N-H bending vibrations. Area 6 is a complex one ($1600-1760\text{ cm}^{-1}$), corresponding to bending vibrations N-H (amino acids), C=O stretchings (aldehydes and cetones, esters) as well to free fatty acids (1710 cm^{-1}) and glycerides (1740 cm^{-1}). Area 7 ($2800-2900\text{ cm}^{-1}$), corresponds to C-H stretching vibrations specific to CH_3 and CH_2 from lipids, metoxy derivatives, C-H (aldehydes), including *cis* double bonds. Area 8 ($3350-3600\text{ cm}^{-1}$) corresponds to stretching vibrations of OH groups (from water, alcohols, phenols, carbohydrates, peroxides) as well from amides (3650 cm^{-1}).

In alcoholic extracts there are absorption peaks in the domain $1300-1800\text{ cm}^{-1}$, more than in glycerin, e.g. at 1558 , 1517 and 1467 cm^{-1} , as well in the region $1380-1450\text{ cm}^{-1}$. Such differences were noticed also by other authors, after processing the second derivative (Liu si col, 2006) in Angelica extracts, where typical signals specific to cellulose and hemicelluloses at 3413 and 1054 cm^{-1} . The signals at 1642 and 1536 cm^{-1} correspond to amide I band (carbonyl group) and amide II (stretching νCN + bending δNH) found in glycoproteins. Carbonyl groups have specific signals at 1743 cm^{-1} . Fig. 4 represents the differences

recorded between the IR areas 1, 2, 4, 6, 7 corresponding to each plant extract (E, G or M).

Looking to region 1 (specific to terpenoids) it has been noticed that plants 6, 5 and 4 had higher peak areas in ethanol, similarly to results from UV-spectra. In the other IR regions (4 and 6) no significant differences between the three solvents extracts were noticed, but in regions 2 (corresponding to glucosides) and 7 (lipids), in all plant extracts, the M extract was significantly more charged in molecules that E or G extracts.

Finally it has been compared the phenolic concentrations, determined by FTIR method, based on the peak intensity at 1743 cm^{-1} and total phenolics calculated from Vis spectrometry. A significant ($p < 0.05$) correlation factor was obtained, as shown in Fig. 5. It is known that the measurement by Vis spectrometry is not specific to phenolics and can overestimate concentrations, while the FTIR method using the peak area ($950-1900\text{ cm}^{-1}$) estimation can give also false results. It can be consider in this case that measurements based on FTIR absorption intensity at 1743 cm^{-1} offer the best evaluation of the phenolics concentration in these plants. Generally the concentrations oscillated between 15 to 20 mg GA/ml methanolic extract (15000-20000 ppm), in agreement with other determinations.

Conclusions

The data presented in this study showed that UV-Vis spectrometry and FT-MIR spectroscopy are adequate techniques to fingerprint comparatively and to evaluate the extraction yield of medicinal herbs with hepatoprotective potential. Based on UV spectrometry, the extraction yields were superior in acidic methanol comparative to glycerin and ethanol, increased in phenolic acids comparative to flavonoid derivatives. Based on the differences of polarity between the three solvents used, higher extraction yields were obtained for dandelion, artichoke, celandine and St. John wort, more rich in phenolic acids than flavonoids. Mary thistle and Wolf's claw had lower concentrations of phenolics, but higher content of lignans and terpenoids. Based on the FT-MIR spectroscopy, for each plant extract it was determined the fingerprint region




Fig. 5. Comparative representation of the total phenolics (expressed as mg GA/ml methanolic extract), calculated from Vis spectrometry, from FTIR peak intensity at 1743 cm^{-1} and from FTIR peak area

to be located between 900 and 1500 cm^{-1} and it has been identified the specific functional groups.

All FTIR data will be correlated and further validated with the detailed HPLC analysis of the same extracts, in order to validate the FTIR method as a good tool to investigate the fingerprint and to predict the composition of medicinal plants or to evaluate the quality and authenticity of different standardized formulas.

Acknowledgements

This work was partly supported from the ANCS-PNII projects AMPRENTAL (ID 51049) and PHYTOMED (ID 61008) (2007-2010).

References

- Ahsan R, Islam M, Bulbul IJ, Musaddik A, Haque E (2009). Hepatoprotective Activity of Methanol Extract of Some Medicinal Plants Against Carbon Tetrachloride-Induced Hepatotoxicity in Rats. *Europ J Sci Res* 37:302-310.
- Ali SM, Olivo M (2002). Bio-distribution and subcellular localization of hypericin and its role in PDT induced apoptosis in cancer cells. *Int J Oncol* 21:531-540.
- Baer-Dubowska W, Szafer H, Krajka-Kuzniak V (1998). Inhibition of murine hepatic cytochrome P450 activities by natural and synthetic phenolic compounds. *Xenobio* 28:735-743.
- Barnes J, Anderson LA, Philippson JD (2001). St. John's wort (*Hypericum perforatum* L.): a review of its chemistry, pharmacology and clinical properties. *J Pharmacol* 53(5):583-600.
- Basaga H, Poli G, Tekkaya C, Aras I (1997). Free radical scavenging and antioxidative properties of 'silibin' complexes on microsomal lipid peroxidation. *Cell Biochem Funct* 15:27-33.
- Bender DA (2005). The promise of metabolomics. *J Sci Food Agric* 85:7-9.
- Benninger J, Schneider HT, Schuppan D, Kirchner T, Hahn EG (1999). Acute hepatitis induced by greater celandine (*Chelidonium majus*). *Gastroenterol* 117:1234-7.
- Betancor-Fernandez A, Pérez-Gálvez A, Sies H, Stahl W (2003). Screening pharmaceutical preparations containing extracts of turmeric rhizome, artichoke leaf, devil's claw root and garlic or salmon oil for antioxidant capacity. *J Pharm Pharmacol* 55(7):981-6.
- Bone K (1996). *Chelidonium*-A medicinal poppy. *MediHerb Professional Newsletter* 49:1-3.
- Bosisio E, Benelli C, Pirola O (1992) Effect of the flavanolignans of *Silybum marianum* L. on lipid peroxidation in rat liver microsomes and freshly isolated hepatocytes. *Pharmacol Res* 25:147-154.
- Cabrera C (1996). Milk Thistle: A clinician's report. *Medical Herbalism* 6:1-5.
- Duke JA (1985). *CRC Handbook of Medicinal Herbs*. CRC Press, Boca Raton, FL.
- Gong F, Bo-Tang W, Yi-Zeng L, Foo-Tim C, Ying-Sing F (2006). Variable selection for discriminating herbal medicines with chromatographic fingerprints. *Analytica Chim. Acta* 572:265-271.
- Gong F, Zhang Q, Zhang J (2009). Quality Assessment of Herbal Medicine with Chromatographic Fingerprint. *J Drug Add, Educ Eradic* 4:257-302.
- Fan XH, Cheng YY, Ye ZL, Lin RC, Qian ZZ (2006). Multiple chromatographic fingerprinting and its application to the quality control of herbal medicines. *Anal Chem Acta* 555:217-224.
- Favari L, Perez-Alvarez V (1997). Comparative effects of colchicine and silymarin on CCl_4 -chronic liver damage in rats. *Arch Med Res* 28:11-17.
- Fehér I, Deak G, Muzes G (1989). Liver-protective action of silymarin therapy in chronic alcoholic liver diseases. *Orv Hetil* 130:2723-2727.
- Ferenci P, Dragosics B, Dittrich H, Frank H, Benda L, Lochs H, Meryn S, Base W, Schneider B (1989). Randomized controlled trial of silymarin treatment in patients with cirrhosis of the liver. *J Hepatol* 9:105-113.
- Gilca M, Gaman L, Panait E, Stoian I, Atanasiu V (2010). Chelidonium majus-an integrative review: traditional knowledge versus modern findings. *Forsch Komplementmed* 17(5):241-248.
- Giri L, Andola HC, Purohit VK, Rawat MSM, Rawal RS, Bhatt ID (2010). Chromatographic and spectral fingerprinting standardization of traditional medicines: an overview as modern tools. *Res J Phytochem* 4:234-241.
- Gu Y, Qian D, Duan J, Wang Z, Guo J, Tang Y, Guo S (2010). Simultaneous determination of seven main alkaloids of *Chelidonium majus* L. by ultra-performance LC with photodiode-array detection. *J Sep Sci* 33:1004-1009.
- Hashimoto A, Kameoka T (2008). Applications of infrared spectroscopy to biochemical, food, and agricultural processes. *App Spectrosc Rev* 43:416-451.
- Hrisu A, Galesanu MR, Moisa L (1980). Cholecystokinetic action of an alkaloid extract of *Chelidonium majus*. *Rev Med Chir Soc Med Nat Iasi* 84:559-561.
- Hussain K, Majeed MT, Ismail Z, Sadikun A, Ibrahim P (2009). Complementary and alternative medicine: quality assessment strategies and safe usage. *Southern Med Rev* 1(2):19-23.
- Hussain K, Ismail Z, Sadikun A, Ibrahim P (2009a). Evaluation of Metabolic Changes in Fruit of Piper Sarmentosum in various Seasons by Metabolomics using Fourier Transform Infrared (FTIR) Spectroscopy. *J Pharm Clin Res* 1(2):68-71.
- Istudor V (2003). Farmacognozie, Fitochimie, Fitoterapie. *Farmacia* 40(2):55-60.
- Jeon HJ, Kang HJ, Jung HJ, Kang YS, Lim CJ, Kim YM, Park EH (2008). Antiinflammatory activity of *Taraxacum officinale*. *J Ethnopharmacol* 115(1):82-88.
- Kurth H, Spreeman R (1998). Phytochemical characterization of various St. John Wort extracts. *Adv Ther* 15:117-128.
- Li S, Han Q, Qiao C, Song J, Cheng CL, Xu H (2008). Chemical

- markers for the quality control of herbal medicines: an overview. *Chinese Med* 3(7):1-18.
- Li H, Xia N, Brausch I, Yao Y, Förstermann U (2004a). Flavonoids from artichoke (*Cynara scolymus* L.) up-regulate endothelial-type nitric-oxide synthase gene expression in human endothelial cells. *J Pharmacol Exp Ther* 310(3):926-932.
- Li Y, Sun S, Zhou Q, Qin Z, Tao J, Wang J, Fang X (2004b). Identification of American ginseng from different region using FT-IR and two-dimensional correlation IR spectroscopy. *Vibrat Spectroscopy* 36:227-232.
- Liang YZ, Xie P, Chan K (2004). Review: Quality control of herbal medicines. *J Chromatography B* 812:53-70.
- Liu H, Sun S, Lv G, Chan KKC (2006). Study on Angelica and its different extracts by Fourier transform infrared spectroscopy and two-dimensional correlation IR spectroscopy. *Spectrochim Acta Part A* 64:321-326.
- Lopus M, Panda D (2006). The benzophenanthridine alkaloid sanguinarine perturbs microtubule assembly dynamics through tubulin binding. A possible mechanism for its antiproliferative activity. *FEBS J* 273:2139-2150.
- Malikova J, Zdarilova A, Hlobilkova A (2006). Effects of sanguinarine and chelerythrine on the cell cycle and apoptosis. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub* 150:5-12.
- Maloney V (2004). Plant metabolomics. *BioTeach J* 4:92-99.
- Marakis G, Walker AF, Middleton RW, Booth JC, Wright J, Pike DJ (2002). Artichoke leaf extract reduces mild dyspepsia in an open study. *Phytomedicine* 9(8):694-699.
- Maros T, Rácz G, Katonai B, Kovács VV (1966). Effects of *Cynara scolymus* extracts on the regeneration of rat liver. 1. *Arzneimittelforschung* 16(2):127-129.
- Mattoli L, Cangi F, Maidecchi A, Ghiara C, Ragazzi E, Tubaro M, Stella L, Tisato F, Traldi P (2006). Metabolomic fingerprinting of plant extracts. *J Mass Spectrometry* 41(12):1534-1545.
- McGuffin M, Hobbs C, Upton R, Goldberg A (1997). American Herbal Products Association's Botanical Safety Handbook. CRC Press, Boca Raton, FL.
- McPartland JM (1996). Viral hepatitis treated with *Phyllanthus amarus* and milk thistle (*Silybum marianum*): A case report. *Complement Med Internat*: 4(2):40-42.
- Miguez MP, Anundi I, Sainz-Pardo LA (1994). Hepatoprotective mechanism of silymarin: no evidence for involvement of cytochrome P4502E1. *Chem Biol Interact* 91:51-63.
- Mitra S, Gole K, Samajdar K, Sur RK, Chakraborty BN (1992). Antihepatotoxic activity of *Chelidonium majus*. *Int J Pharmacognosy* 30:125-128.
- Mitra S, Sur RK, Roy A, Mukherjee AS (1996). Effect of *Chelidonium majus* L on experimental hepatic tissue injury. *Phytother Res* 10:354-356.
- Muriel P, Garcipina T, Perez-Alvarez V, Mourelle M (1992). Silymarin protects against paracetamol induced lipid peroxidation and liver damage. *J Appl Toxicol* 12:439-442.
- Nahrstedt A, Butterweck V (1997). Biologically active and other chemical constituents of the herb of *Hypericum perforatum* L. *Pharmacopsychiat* 30:129-134.
- Negi AS, Kumar JK, Luqman S, Shanker K, Gupta MM, Khanuja SP (2008). Recent advances in plant hepatoprotectives: a chemical and biological profile of some important leads. *Med Res Rev* 28(5):746-772.
- Perez-Garcia F, Adzet T, Cañigüeral S (2000). Activity of artichoke leaf extract on reactive oxygen in human leukocytes. *Free Rad Res* 33(5):661-665.
- Ritter R, Schatton WFH (1993). Clinical trial on standardized celandine extract in patients with functional epigastric complaints: Results of placebo-controlled double-blind trial. *Comp Ther Med* 1:189-193.
- Saénz Rodríguez T, García Giménez D, de la Puerta Vázquez R (2002). Choleretic activity and biliary elimination of lipids and bile acids induced by an artichoke leaf extract in rats. *Phytomedicine* 9(8):687-693.
- Salmi HA, Sarna S (1982). Effect of silymarin on chemical, functional, and morphological alterations of the liver; a double blind controlled study. *Scand J Gastroenterol* 17:517-521.
- Schultz H, Baranska M (2007). Identification and quantification of valuable substances by IR and Raman spectroscopy. *Vibrational Spectroscopy* 43:13-25.
- Schütz K, Muks E, Carle R, Schieber A (2006). Separation and quantification of inulin in selected artichoke (*Cynara scolymus* L.) cultivars and dandelion (*Taraxacum officinale* WEB. ex WIGG.) roots by high-performance anion exchange chromatography with pulsed amperometric detection. *Biomed Chromatography* 20(12):1295-1303.
- Singleton VL (1999). Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Methods Enzymol* 299:152-178.
- Utrilla MP (1996). Natural products with hepatoprotective action. *Methods Find Exp Clin Pharmacol* 18(B):11-12.
- Vavreckova C, Gawlik I, Muller K (1996). Benzophenanthridine alkaloids of *Chelidonium majus*; I. Inhibition of 5- and 12-lipoxygenase by a non-redox mechanism. *Planta Med* 62:397-401.
- Wang MF, Simon JE, Aviles IF, He K, Zheng QY, Tadmor Y (2003). Analysis of antioxidative phenolic compounds in artichoke *Cynara scolymus* L. *J Agr Food Chem* 51:601-608.
- WHO (2003). WHO guidelines on Good Agriculture and Collection Practices (GACP) for medicinal plants. Geneva.
- Williams CA, Goldstone F, Greenham J (1996). Flavonoids, cinnamic acids and coumarins from the different tissues and medicinal preparations of *Taraxacum officinale*. *Phytochem* 42(1):121-127.
- Yadav NP, Dixit VK (2008). Recent approaches in herbal drug standardization. *Int J Integr Biol* 2:195-203.
- Zdařilová A, Malikova J, Dvorak Z, Ulrichova J, Simanek V (2006). Quaternary isoquinoline alkaloids sanguinarine and chelerythrine. *In vitro* and *in vivo* effects. *Chemicke Listy* 100:30-41.